

University of Groningen

## BLOCK KRONECKER PRODUCTS AND THE VECB OPERATOR

Koning, Ruud H.; Neudecker, H.; Wansbeek, T.J.

*Published in:*  
Linear Algebra and Its Applications

*DOI:*  
[10.1016/0024-3795\(91\)90332-Q](https://doi.org/10.1016/0024-3795(91)90332-Q)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1991

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Koning, R. H., Neudecker, H., & Wansbeek, T. J. (1991). BLOCK KRONECKER PRODUCTS AND THE VECB OPERATOR. *Linear Algebra and Its Applications*, 149, 165-184. [https://doi.org/10.1016/0024-3795\(91\)90332-Q](https://doi.org/10.1016/0024-3795(91)90332-Q)

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

✓

MEMORANDUM

from

INSTITUTE OF ECONOMIC RESEARCH

FACULTY OF ECONOMICS

UNIVERSITY OF GRONINGEN

P.O. Box 800

9700 AV Groningen

Research Memorandum nr. 351

BLOCK KRONECKER PRODUCTS AND THE VECB OPERATOR

by

Ruud H. Koning  
Heinz Neudecker  
Tom Wansbeek

January 1990

Editors:

Prof.dr. J.L. Bouma  
Prof.dr. S.K. Kuipers  
Prof.dr. P.S.H. Leeflang  
Prof.dr. A. Maddison  
Prof.dr. J. Pen  
Prof.dr. J. Ponstein  
Prof.dr. H-J. Wagener  
Prof.dr. T.J. Wansbeek

TC 1286  
wnee  
5006  
002  
(351)







## BLOCK KRONECKER PRODUCTS AND THE VECB OPERATOR

Ruud H. Koning<sup>\*</sup>

Heinz Neudecker<sup>\*\*</sup>

Tom Wansbeek<sup>\*</sup>

<sup>\*</sup> Department of Economics, University of Groningen, P.O. Box 800, 9700 AV, Groningen, The Netherlands.

<sup>\*\*</sup> Department of Actuarial Sciences and Econometrics, University of Amsterdam, Jodenbreestraat 23, 1011 NH, Amsterdam, The Netherlands.

This work was started when the second author was at the Indian Statistical Institute, New Delhi.







## INTRODUCTION

Almost twenty years ago Singh [7] and Tracy and Singh [9] introduced a generalization of the Kronecker product  $A \otimes B$ . They used varying notation for this new product, viz.  $A \pi B$  and  $A \square B$ . Recently, Hyland and Collins [1] studied the same product under rather restrictive order conditions. They denoted it  $A \bar{\otimes} B$ . Curiously enough, they were not aware of their predecessors' work.

Tracy and Jinadasa [8] considered the product again very recently. They elected to use the notation  $A \pi B$ .

Closely related to the new product is a generalization of the vec operator, which Hyland and Collins very aptly named 'vecb' operator. This and another very similar operator can also be found in the work of the other authors. Hyland and Collins integrated the new product and the vecb operator in a nice way.

In the present paper we define a less restricted generalized Kronecker product. Whereas the product studied before is based on the partitioning of both matrices  $A$  and  $B$ , ours presupposes a partition of  $B$  solely. Researchers can partition  $A$  according to their needs and wishes. This approach is mathematically less cumbersome and very suitable for certain unbalanced partitionings. We propose to use the notation  $A \square B$  for this generalized Kronecker product. The other product will be denoted  $A \pi B$ . Both products will be called 'block Kronecker products'.

The term 'vecb' operator will be used to cover the two variants of the generalized vec operator. These will be distinguished notationally as  $\text{vecb}_c$  and  $\text{vecb}_r$ , and will be shown to be compatible with the two above mentioned block Kronecker products.

In the first section of this paper we discuss the case of balanced partitioning. This involves the product  $A \pi B$ , the  $\text{vecb}_c$  operator and the so-called tilde transform.

In the second section we elaborate the general case of unbalanced partitioning, where we introduce the block Kronecker product  $A \square B$  and the compatible  $\text{vecb}_r$  operator.

In the third section we consider linear equations in partitioned matrices.

Finally, we apply some of the techniques discussed in the fourth and last section.



# 1 BALANCED PARTITIONING

## 1.1 The $\text{vecb}_c$ operator, the tilde transform and the balanced block Kronecker product $A \pi B$

Throughout section 1 the matrix  $A$  is a blockwise partitioned matrix,  $A = (A_{ij})$ , where  $A_{ij}$  is of order  $m \times n$  ( $i = 1, \dots, p; j = 1, \dots, q$ ). So  $A$  is of order  $mp \times nq$ , and we can write

$$A = \sum_{ij} (E_{ij} \otimes A_{ij}), \quad (1)$$

where the matrix  $E_{ij}$  is of order  $p \times q$ , having a unit element in the  $(i, j)$ th position and zeros elsewhere. Obviously, the order of  $E_{ij}$  corresponds to the partition of  $A$ . We define

$$\text{vecb}_c A \equiv \sum_{ij} (\text{vec} E_{ij} \otimes \text{vec} A_{ij}). \quad (2)$$

Thus, if  $p = q = 2$ , so

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix},$$

then

$$\text{vecb}_c A = \begin{bmatrix} \text{vec} A_{11} \\ \text{vec} A_{21} \\ \text{vec} A_{12} \\ \text{vec} A_{22} \end{bmatrix}.$$

We have vectorized the partitioned matrix  $A$  column-blockwise. This is only one way of vectorizing the partitioned matrix  $A$ , in agreement with the literature (Tracy and Singh [9], Singh [7], Tracy and Jinadasa [8], Hyland and Collins [1]). However, we can also vectorize row-blockwise, i.e., vectorize each row of blocks, and stack the resulting vectors. This leads to another type of  $\text{vecb}$  operator:

$$\text{vecb}_r A = \sum_{ij} (\text{vec} E_{ji} \otimes \text{vec} A_{ij}) \quad (3)$$

and for the example we have



$$\text{vecb}_r A = \begin{bmatrix} \text{vec} A_{11} \\ \text{vec} A_{12} \\ \text{vec} A_{21} \\ \text{vec} A_{22} \end{bmatrix}.$$

For the rest of this section we only consider the  $\text{vecb}_c$  operator. We consider the  $\text{vecb}_r$  operator again in section 2, where we discuss the case of unbalanced partitioning.

If we denote the commutation matrix (see Magnus and Neudecker [4]) by  $K$ , then the following property is easily established:

$$\text{vecb}_c A = (I_q \otimes K_{pn} \otimes I_m) \text{vec} A, \quad (4)$$

since

$$\begin{aligned} \text{vecb}_c A &= \sum_{ij} (\text{vec} E_{ij} \otimes \text{vec} A_{ij}) = \sum_{ij} (I_q \otimes K_{pn} \otimes I_m) \text{vec} (E_{ij} \otimes A_{ij}) \\ &= (I_q \otimes K_{pn} \otimes I_m) \sum_{ij} \text{vec} (E_{ij} \otimes A_{ij}) = (I_q \otimes K_{pn} \otimes I_m) \text{vec} A, \end{aligned}$$

by virtue of the result

$$\text{vec} (Y \otimes Z) = (I_q \otimes K_{np} \otimes I_m) (\text{vec} Y \otimes \text{vec} Z) \quad (5)$$

for matrices  $Y$  ( $p \times q$ ) and  $Z$  ( $m \times n$ ) (see Neudecker and Wansbeek [6]).

Closely connected with this is the so-called tilde transform, which links the  $\text{vec}$  operator and the  $\text{vecb}_c$  operator. The tilde transform of  $A$  is defined by

$$\tilde{A} = \sum_{ij} (\text{vec} A_{ij}) (\text{vec} E_{ij})'. \quad (6)$$

For example, if again  $p = q = 2$ , we get

$$\tilde{A} = [\text{vec} A_{11}, \text{vec} A_{21}, \text{vec} A_{12}, \text{vec} A_{22}].$$

From (4), (5) and (6) we have

$$\text{vec} \tilde{A} = \text{vecb}_c A = (I_q \otimes K_{pn} \otimes I_m) \text{vec} A. \quad (7)$$

We next consider the equation

$$\text{vecb}_c B C A' = X \text{vecb}_c C, \quad (8)$$



for  $A$  as defined above,

$$B = (B_{kl}) = \sum_{kl} (E_{kl} \otimes B_{kl}),$$

with  $B_{kl}$  of order  $u \times v$  ( $k = 1, \dots, s; l = 1, \dots, t$ ) and

$$C = (C_{lj}) = \sum_{lj} (E_{lj} \otimes C_{lj}),$$

with  $C_{lj}$  of order  $v \times n$  ( $l = 1, \dots, t; j = 1, \dots, q$ ). Using (4) we have

$$\begin{aligned} \text{vecb}_c B C A' &= (I_p \otimes K_{sm} \otimes I_u) \text{vec} B C A' = (I_p \otimes K_{sm} \otimes I_u) (A \otimes B) \text{vec} C \\ &= (I_p \otimes K_{sm} \otimes I_u) (A \otimes B) (I_q \otimes K_{nt} \otimes I_v) \text{vecb}_c C, \end{aligned} \quad (9)$$

from which follows the solution

$$\text{vecb}_c B C A' = (A \pi B) \text{vecb}_c C, \quad (10)$$

with

$$A \pi B \equiv (I_p \otimes K_{sm} \otimes I_u) (A \otimes B) (I_q \otimes K_{nt} \otimes I_v). \quad (11)$$

This defines the balanced block Kronecker product  $A \pi B$  (cf. Tracy and Singh [9] and Hyland and Collins [1], formula (2.4)). If  $p = q = s = 2$  and  $t = 3$ , the block Kronecker product of  $A$  and  $B$  takes the form

$$A \pi B = \begin{bmatrix} A_{11} \otimes B_{11} & A_{11} \otimes B_{12} & A_{11} \otimes B_{13} & A_{12} \otimes B_{11} & A_{12} \otimes B_{12} & A_{12} \otimes B_{13} \\ A_{11} \otimes B_{21} & A_{11} \otimes B_{22} & A_{11} \otimes B_{23} & A_{12} \otimes B_{21} & A_{12} \otimes B_{22} & A_{12} \otimes B_{23} \\ A_{21} \otimes B_{11} & A_{21} \otimes B_{12} & A_{21} \otimes B_{13} & A_{22} \otimes B_{11} & A_{22} \otimes B_{12} & A_{22} \otimes B_{13} \\ A_{21} \otimes B_{21} & A_{21} \otimes B_{22} & A_{21} \otimes B_{23} & A_{22} \otimes B_{21} & A_{22} \otimes B_{22} & A_{22} \otimes B_{23} \end{bmatrix}.$$

If  $A$  is partitioned elementwise (i.e.,  $m = n = 1$ ), the  $\text{vecb}_c$  operator and the block Kronecker product reduce to the usual  $\text{vec}$  operator and Kronecker product.

This derivation of the balanced block Kronecker product  $A \pi B$  is inspired by Hyland and Collins. We prefer it to the explicit definitions of Tracy and Singh and Singh and Jinadasa, since this derivation shows how the product arises in a natural manner from vectorizing a partitioned matrix. We can write  $A \pi B$  in a form that elegantly generalizes (1):



$$A\pi B = \sum_{ijkl} \left\{ (E_{ij} \otimes E_{kl}) \otimes (A_{ij} \otimes B_{kl}) \right\}, \quad (11')$$

because

$$\begin{aligned} A\pi B &= (I_p \otimes K_{sm} \otimes I_u) (A \otimes B) (I_q \otimes K_{nt} \otimes I_v) \\ &= \sum_{ijkl} (I_p \otimes K_{sm} \otimes I_u) (E_{ij} \otimes A_{ij} \otimes E_{kl} \otimes B_{kl}) (I_q \otimes K_{nt} \otimes I_v) \\ &= \sum_{ijkl} \left\{ E_{ij} \otimes K_{sm} (A_{ij} \otimes E_{kl}) K_{nt} \otimes B_{kl} \right\} \\ &= \sum_{ijkl} \left\{ (E_{ij} \otimes E_{kl}) \otimes (A_{ij} \otimes B_{kl}) \right\}. \end{aligned}$$

Here the orders of the matrices  $E_{ij}$  and  $E_{kl}$  correspond to the respective partitions of  $A$  and  $B$ . Hence  $E_{ij}$  is of order  $p \times q$  and  $E_{kl}$  is of order  $s \times t$ .

## 1.2 Properties of the $\text{vecb}_c$ operator and the balanced block Kronecker product $A\pi B$

1. Let  $a = (a_i)$  and  $b = (b_k)$ , where  $a_i$  is of order  $m \times 1$  and  $b_k$  is of order  $u \times 1$  ( $i = 1, \dots, p$ ;  $k = 1, \dots, s$ ), then

$$a\pi b = \text{vecb}_c b a' = (I_p \otimes K_{sm} \otimes I_u) (a \otimes b). \quad (12)$$

Proof: Replace  $A$  by  $a$ ,  $B$  by  $b$  and  $n, q, v, t$  by 1 in (11). Replace  $C$  by 1 in (10).

2. Let  $A$  and  $B$  be defined as above. Then

$$\text{vecb}_c (A\pi B) = (I_q \otimes K_{tp} \otimes I_{sn} \otimes K_{vm} \otimes I_u) (\text{vecb}_c A \pi \text{vecb}_c B) \quad (13)$$

Proof:

$$\begin{aligned} \text{vecb}_c A \pi \text{vecb}_c B &= (I_{pq} \otimes K_{st, mn} \otimes I_{uv}) (\text{vecb}_c A \otimes \text{vecb}_c B) \\ &= (I_{pq} \otimes K_{st, mn} \otimes I_{uv}) \sum_{ijkl} (\text{vec} E_{ij} \otimes \text{vec} A_{ij} \otimes \text{vec} E_{kl} \otimes \text{vec} B_{kl}) \\ &= \sum_{ijkl} (\text{vec} E_{ij} \otimes \text{vec} E_{kl} \otimes \text{vec} A_{ij} \otimes \text{vec} B_{kl}), \end{aligned}$$

by virtue of (2) and (11); furthermore,

$$I_q \otimes K_{tp} \otimes I_{sn} \otimes K_{vm} \otimes I_u = (I_q \otimes K_{tp} \otimes I_s) \otimes (I_n \otimes K_{vm} \otimes I_u),$$



hence we find

$$\begin{aligned}
& (I_q \otimes K_{tp} \otimes I_{sn} \otimes K_{vm} \otimes I_u) (\text{vecb}_c A \pi \text{vecb}_c B) \\
&= \sum_{ijkl} \left\{ \text{vec}(E_{ij} \otimes E_{kl}) \otimes \text{vec}(A_{ij} \otimes B_{kl}) \right\} \\
&= \text{vecb}_c \sum_{ijkl} \left\{ (E_{ij} \otimes E_{kl}) \otimes (A_{ij} \otimes B_{kl}) \right\} \\
&= \text{vecb}_c (A \pi B),
\end{aligned}$$

by using (2) and (11').

3. When  $A$  is elementwise partitioned ( $m = n = 1$ ), then

$$\text{vecb}_c(A \otimes B) = \text{vec} A \otimes \text{vec} B. \quad (14)$$

If further  $C = A \otimes B$ , then

$$\tilde{C} = (\text{vec} B) (\text{vec} A)'. \quad (15)$$

Proof:

$$A = \sum_{ij} a_{ij} E_{ij},$$

hence

$$\begin{aligned}
\text{vecb}_c(A \otimes B) &= \sum_{ij} a_{ij} \text{vecb}_c(E_{ij} \otimes B) = \sum_{ij} a_{ij} (\text{vec} E_{ij} \otimes \text{vec} B) = \\
&= \text{vec} A \otimes \text{vec} B
\end{aligned}$$

by (2).

Further,

$$\begin{aligned}
\tilde{C} &= \sum_{ij} (\text{vec} C_{ij}) (\text{vec} E_{ij})' = \sum_{ij} (\text{vec} a_{ij} B) (\text{vec} E_{ij})' \\
&= (\text{vec} B) \left( \sum_{ij} \text{vec} a_{ij} E_{ij} \right)' = (\text{vec} B) (\text{vec} A)'.
\end{aligned}$$

4. When  $A$  and  $B$  are columnwise partitioned ( $n = v = 1$ ), then

$$\text{vecb}_c \{ (\text{vec} B) (\text{vec} A)' \} = \text{vec} (A \otimes B). \quad (16)$$



If further  $C = (\text{vec}B)(\text{vec}A)'$ , then

$$\hat{C} = A \otimes B. \quad (17)$$

Proof:

$$\text{vec}_c \{ (\text{vec}B)(\text{vec}A)' \} = (I_q \otimes K_{t,mp} \otimes I_{us}) (\text{vec}A \otimes \text{vec}B) = \text{vec}A \otimes B,$$

by virtue of (12) and (5). Finally,

$$\begin{aligned} C &= \sum_{ij} \{ E_{ij} \otimes (B_{.i} A'_{.j}) \} \\ \hat{C} &= \sum_{ij} \{ [\text{vec}(B_{.i} A'_{.j})] [\text{vec}E_{ij}]' \} \\ &= \sum_{ij} (A_{.j} \otimes B_{.i}) (e'_j \otimes e'_i) \\ &= \left[ \sum_j A_{.j} e'_j \right] \otimes \left[ \sum_i B_{.i} e'_i \right] \\ &= A \otimes B, \end{aligned}$$

by (1) and (6).

5. Let  $A, B, C, D$  be partitioned matrices of order  $mp \times nq, nq \times rk, rk \times sl$  and  $mp \times sl$  whose blocks are of order  $m \times n, n \times k, k \times l$  and  $l \times m$ , respectively. Then

$$\text{tr}ABCD' = (\text{vec}A')' (D \otimes B) \text{vec}C = (\text{vec}_c A')' (D \pi B) \text{vec}_c C \quad (18)$$

Proof:

$$\begin{aligned} \text{tr}ABCD' &= (\text{vec}A')' (D \otimes B) \text{vec}C \\ &= (\text{vec}A')' (I_p \otimes K_{mq} \otimes I_n) (I_p \otimes K_{qm} \otimes I_n) (D \otimes B) \\ &\quad (I_s \otimes K_{lr} \otimes I_k) (I_s \otimes K_{rl} \otimes I_k) \text{vec}C \\ &= (\text{vec}_c A')' (D \pi B) \text{vec}_c C, \end{aligned}$$

by virtue of (11).

6. When  $A$  and  $B$  are columnwise partitioned ( $n = v = 1$ ),



$$A\pi B = A\otimes B. \quad (19)$$

Proof:

$$A\pi B = [I_1 \otimes K_{1,mp} \otimes I_{us}] (A \otimes B) [I_q \otimes K_{1t} \otimes I_1] = A \otimes B,$$

by (11).

In section 2.2 we prove sixteen properties of the block Kronecker product in the unbalanced case. *Mutatis mutandis*, the same properties hold for the case of balanced partitioning as discussed here.

## 2 UNBALANCED PARTITIONING

### 2.1 The $\text{vecb}_r$ operator and the block Kronecker product

Let us consider the blockwise partitioned matrix

$$A = \begin{bmatrix} A_{11} & \dots & A_{1q} \\ \vdots & & \vdots \\ A_{p1} & \dots & A_{pq} \end{bmatrix} \equiv \begin{bmatrix} A_{1.} \\ \vdots \\ A_{p.} \end{bmatrix},$$

with typical block  $A_{ij}$  of order  $m_i \times n_j$ . Furthermore,  $m \equiv \sum_{i=1}^p m_i$  and  $n \equiv \sum_{j=1}^q n_j$ . So contrary to what we had above,  $A$  is now an  $m \times n$ -matrix, not all submatrices being of the same size. This is an example of a matrix partitioned in an inbalanced way. Various authors have defined the  $\text{vecb}_c$  operator on this matrix, which they call  $\text{vecb}$ .

Let us consider the equation

$$\text{vecb}_c B C A' = X \text{vecb}_c C,$$

with  $B$  a  $u \times v$  matrix consisting of blocks  $B_{kl}$  of order  $u_k \times v_l$  ( $u = \sum_{k=1}^s u_k$ ,  $v = \sum_{l=1}^r v_l$ ) and  $C$  an  $v \times n$  matrix consisting of blocks  $v_l \times n_j$ . The solution is then the block Kronecker product  $A\pi B$ . We have

$$(B C A')_{ki} = \sum_{jl} (B_{kl} C_{lj} A'_{ji})$$

$$\text{vec}(B C A')_{ki} = \sum_{jl} \text{vec}(B_{kl} C_{lj} A'_{ji}) = \sum_{jl} (A_{ij} \otimes B_{kl}) \text{vec} C_{lj}$$



$$= [A_{i1} \otimes B_{k1}, \dots, A_{i1} \otimes B_{kt}, \dots, A_{iq} \otimes B_{k1}, \dots, A_{iq} \otimes B_{kt}] \text{vecb}_c C.$$

Hence

$$\begin{aligned} \text{vecb}_c BCA' &= \begin{bmatrix} A_{11} \otimes B_{11} & \dots & A_{11} \otimes B_{1t} & \dots & A_{1q} \otimes B_{11} & \dots & A_{1q} \otimes B_{1t} \\ A_{11} \otimes B_{s1} & \dots & A_{11} \otimes B_{st} & \dots & A_{1q} \otimes B_{s1} & \dots & A_{1q} \otimes B_{st} \\ \vdots & & \vdots & & \vdots & & \vdots \\ A_{p1} \otimes B_{11} & \dots & A_{p1} \otimes B_{1t} & \dots & A_{pq} \otimes B_{11} & \dots & A_{pq} \otimes B_{1t} \\ A_{p1} \otimes B_{s1} & \dots & A_{p1} \otimes B_{st} & \dots & A_{pq} \otimes B_{s1} & \dots & A_{pq} \otimes B_{st} \end{bmatrix} \text{vecb}_c C \\ &= (A \pi B) \text{vecb}_c C. \end{aligned}$$

A restrictive feature of this approach is that all three matrices have to be partitioned. However, by vectorizing in a different manner, this problem can be avoided.

Let us therefore consider the equation

$$\text{vecb}_r BCA' = X \text{vecb}_r C, \quad (20)$$

with  $B$  as defined above but  $A$  is a not necessarily partitioned  $m \times n$  matrix.

Using the notation introduced above, we have as the  $k^{\text{th}}$  block row of  $BCA'$

$$(BCA')_k = \sum_l B_{kl} C_l A' \quad k = 1, \dots, s$$

hence,

$$\text{vec}(BCA')_k = \sum_l (A \otimes B_{kl}) \text{vec} C_l = [A \otimes B_{k1}, \dots, A \otimes B_{kt}] \text{vecb}_r C$$

and consequently

$$\text{vecb}_r BCA' = \begin{bmatrix} A \otimes B_{11} & \dots & A \otimes B_{1t} \\ \vdots & & \vdots \\ A \otimes B_{s1} & \dots & A \otimes B_{st} \end{bmatrix} \text{vecb}_r C.$$

The solution turns out to be another block Kronecker product, which we will denote by " $\square$ ":

$$X = A \square B \equiv \begin{bmatrix} A \otimes B_{11} & \dots & A \otimes B_{1t} \\ \vdots & & \vdots \\ A \otimes B_{s1} & \dots & A \otimes B_{st} \end{bmatrix}. \quad (21)$$

Note that we did not partition  $A$ , since there is no mathematical need to do



so. One can partition  $A$  if one wants and as one pleases, given the nature of the practical problem one is concerned with. The fact that it is unnecessary to partition  $A$  follows from our different way of vectorization. While taking the  $\text{vecb}_c$  of a matrix  $A$ , one makes a "jump" from  $A_{11}$  to  $A_{21}$ . The  $\text{vecb}_r$  operator does not make this jump, and in that sense, this operator preserves more of the structure of the matrix  $A$ .

## 2.2 Properties of the block Kronecker product $A \square B$

We shall prove sixteen properties of the block Kronecker product  $A \square B$ , more or less following the set-up of Hyland and Collins. They derive similar properties for the block Kronecker product  $A \pi B$ . We shall, however, impose less restrictive order conditions than they did. Moreover, the properties of the product  $A \square B$  are much easier to derive than for  $A \pi B$ , if the partition is unbalanced. The properties are:

1.  $\text{vecb}_r(BCA') = (A \square B) \text{vecb}_r C.$

Proof: Already supplied in Section 2.1.

2.  $\text{vecb}_r(AD + DB) = (I \square A + B' \square I) \text{vecb}_r D.$

Proof:

$$\begin{aligned} \text{vecb}_r(AD + DB) &= \text{vecb}_r ADI + \text{vecb}_r IDB = (I \square A) \text{vecb}_r D \\ &+ (B' \square I) \text{vecb}_r D = (I \square A + B' \square I) \text{vecb}_r D. \end{aligned}$$

3.  $A \square B = K_1(B \otimes A)K_2$  for some permutation matrices  $K_1$  and  $K_2$ .

Proof: Let  $A$  be an  $m \times n$  matrix and let  $B$  be a  $u \times v$  matrix, with blocks of order  $u_l \times v_k$  ( $l = 1, \dots, r, k = 1, \dots, s$ ). Then

$$B \otimes A = \begin{bmatrix} B_{11} \otimes A & \dots & B_{1s} \otimes A \\ \vdots & & \vdots \\ B_{r1} \otimes A & \dots & B_{rs} \otimes A \end{bmatrix} = \begin{bmatrix} K_{u_1 m} & & \\ & \ddots & \\ & & K_{u_r m} \end{bmatrix} (A \square B) \begin{bmatrix} K_{nv_1} & & \\ & \ddots & \\ & & K_{nv_s} \end{bmatrix}.$$

Hence

$$A \square B = \begin{bmatrix} K_{mu_1} & & \\ & \ddots & \\ & & K_{mu_r} \end{bmatrix} (B \otimes A) \begin{bmatrix} K_{v_1 n} & & \\ & \ddots & \\ & & K_{v_s n} \end{bmatrix}.$$



4.  $(A+B) \square C = A \square C + B \square C.$

Proof: Evident.

5.  $A \square (B+C) = A \square B + A \square C.$

Proof: Evident.

6.  $(A \square B)' = A' \square B'.$

Proof: Follows from 3.

7.  $(A \square B)(C \square D) = AC \square BD.$

Proof:

$$\begin{aligned} (A \square B)(C \square D) \text{vecb}_r X &= (A \square B) \text{vecb}_r D X C' = \text{vecb}_r B D X C' A' \\ &= \text{vecb}_r B D X (A C)' = (A C \otimes B D) \text{vecb}_r X. \end{aligned}$$

As this holds for any  $X$ , the result follows.

8.  $(A \square B)^- = A^- \square B^-$ , for any generalized inverse.

Proof: Follows from 7.

9.  $B \square A = K_1(A \square B)K_2$  for some permutation matrices  $K_1$  and  $K_2$ .

Proof: Follows from 3.

10.  $|A \square B| = |A|^n \cdot |B|^m$  for square matrices  $A$   $m \times m$  and  $B$   $n \times n$ .

Proof: Use 3 to get

$$|A \square B| = |B \otimes A| = |A|^n \cdot |B|^m.$$

Note: The partition is clearly immaterial.

11.  $\text{tr}(A \square B) = \text{tr} A \cdot \text{tr} B$  for square matrices  $A$  and  $B$  and  $B$  with square diagonal blocks.

Proof: Let  $B_{11}, \dots, B_{nn}$  be the square diagonal blocks of  $B$ . Then

$$\text{tr}(A \square B) = \sum_i \text{tr}(A \otimes B_{ii}) = \text{tr} A \cdot (\sum_i \text{tr} B_{ii}) = \text{tr} A \cdot \text{tr} B.$$

12.  $(I \square A)(B \square I) = (B \square I)(I \square A) = B \square A.$

Proof: Follows from 7 applied twice.



13.  $\exp(A \square I + I \square B) = \exp A \square \exp B.$

Proof:

$$\begin{aligned}
 \exp(A \square I + I \square B) &= \exp(A \square I) \exp(I \square B) = \exp\{K(I \otimes A)K'\} \exp\{K(B \otimes I)K'\} \\
 &= \sum_{i=0}^{\infty} \{K(I \otimes A)K'\}^i \cdot \sum_{j=0}^{\infty} \{K(B \otimes I)K'\}^j \\
 &= K \sum_{i=0}^{\infty} (I \otimes A)^i K' \cdot K \sum_{j=0}^{\infty} (B \otimes I)^j K' \\
 &= K \sum_{i=0}^{\infty} (I \otimes A^i) \cdot \sum_{j=0}^{\infty} (B^j \otimes I) K' = K(I \otimes \exp A)(\exp B \otimes I)K' \\
 &= K(\exp B \otimes \exp A)K' = \exp A \square \exp B.
 \end{aligned}$$

14. The matrices  $A \square B$  and  $B \otimes A$  have identical sets of eigenvalues, for square  $A$  and  $B$ .

Proof: Suppose  $A(B)$  has eigenvalues  $\lambda_i(\mu_j)$  and associated eigenvectors  $x_i(y_j)$ . Then

$$(B \otimes A)(y_j \otimes x_i) = \lambda_i \mu_j (y_j \otimes x_i).$$

Hence, by virtue of 3,

$$\begin{aligned}
 K_1(A \square B)K_2(y_j \otimes x_i) &= K_1(A \square B)(x_i \square y_j) = K_1(Ax_i \square By_j) \\
 &= \lambda_i \mu_j K_1(x_i \square y_j),
 \end{aligned}$$

or

$$(A \square B)(x_i \square y_j) = \lambda_i \mu_j (x_i \square y_j).$$

15. The matrices  $A \square I + I \square B$  and  $I \square A + B \square I$  have identical sets of eigenvalues.

Proof: Let  $A$  and  $B$  be defined as above. Then

$$(I \otimes A + B \otimes I)(y_j \otimes x_i) = (\lambda_i + \mu_j) \cdot (y_j \otimes x_i).$$

Hence

$$\begin{aligned}
 K'_1(A \square I + I \square B)K'_2(y_j \otimes x_i) &= K'_1(A \square I + I \square B)(x_i \square y_j) \\
 &= K'_1(Ax_i \square y_j + x_i \square By_j) = (\lambda_i + \mu_j)K'_1(x_i \square y_j),
 \end{aligned}$$

or



$$(A \otimes I + I \otimes B)(x_i \otimes y_j) = (\lambda_i + \mu_j)(x_i \otimes y_j).$$

16. When  $A$  has eigenvectors  $x_i$  and  $B$  has eigenvectors  $y_j$ , then  $A \otimes B$  and  $A \otimes I + I \otimes B$  will have eigenvectors  $x_i \otimes y_j$ .

Proof: Has been supplied under 14 and 15.

As stated above, Hyland and Collins presented identical properties for the  $\text{vecb}_c$  operator. Unfortunately, their block Kronecker product was unnecessarily constrained as to its partitioning. Tracy and Singh proved the properties 4, 5, 11 and 12 without unnecessarily constraining their matrices.

### 3 THE EQUATION $\text{vec}X = \text{vecb}_r C$ .

We shall now consider the equation

$$\text{vec}X = \text{vecb}_r C. \quad (22)$$

When  $C$  is appropriately partitioned we can solve for  $X$ . The solution is not always unique, even if we preserve the partition of  $C$ . Take for example

$$C = \begin{bmatrix} C_1 \\ \vdots \\ C_p \end{bmatrix} \begin{matrix} m_1 \\ \vdots \\ m_p \end{matrix}.$$

Then we have the solutions

$$X_1 = (\text{vec}C_1, \dots, \text{vec}C_p)$$

and

$$X_2 = (C_1, \dots, C_p),$$

because

$$\text{vecb}_r C = \begin{bmatrix} \text{vec}C_1 \\ \vdots \\ \text{vec}C_p \end{bmatrix} = \text{vec}(\text{vec}C_1, \dots, \text{vec}C_p) = \text{vec}(C_1, \dots, C_p).$$



Another example is

$$C = \begin{bmatrix} \overset{n}{C_{11}} \cdot \overset{n}{C_{1q}} \\ \vdots \\ \underset{m}{C_{p1}} \cdot \underset{m}{C_{pq}} \end{bmatrix}_m,$$

where all blocks have the same order  $(m \times n)$ . Then we have one solution

$$X = (\text{vec}C_{11}, \dots, \text{vec}C_{1q}, \dots, \text{vec}C_{p1}, \dots, \text{vec}C_{pq}),$$

because

$$\begin{aligned} \text{vec}_r C &= \begin{bmatrix} \text{vec}C_1 \\ \vdots \\ \text{vec}C_p \end{bmatrix} = \begin{bmatrix} \text{vec}(C_{11}, \dots, C_{1q}) \\ \vdots \\ \text{vec}(C_{p1}, \dots, C_{pq}) \end{bmatrix} \\ &= \begin{bmatrix} \text{vec}C_{11} \\ \text{vec}C_{1q} \\ \text{vec}C_{p1} \\ \text{vec}C_{pq} \end{bmatrix} = \text{vec}(\text{vec}C_{11}, \dots, \text{vec}C_{1q}, \dots, \text{vec}C_{p1}, \dots, \text{vec}C_{pq}). \end{aligned}$$

A third example is

$$C = \begin{bmatrix} c_{11} \cdot c_{1q} \\ \vdots \\ c_{p1} \cdot c_{pq} \end{bmatrix},$$

the matrix being partitioned elementwise.

Now we have the solution

$$X = C',$$

because

$$\text{vec}_r C = (c_{11}, \dots, c_{1q}, \dots, c_{p1}, \dots, c_{pq})' = \text{vec}C'.$$

We have the following two results. First, if  $C = A \otimes B$ , where  $A$  is elementwise partitioned, then

$$X_1 = (\text{vec}B)(\text{vec}A)', \quad (23a)$$

$$X_2 = (\text{vec}A')' \otimes B \quad (23b)$$



are solutions of the equation  $\text{vec}X = \text{vecb}_r C$ . Since

$$C = A \otimes B = \begin{bmatrix} a_{11}B & \dots & a_{1n}B \\ \vdots & & \vdots \\ a_{m1}B & \dots & a_{mn}B \end{bmatrix},$$

say, we have

$$\begin{aligned} \text{vecb}_r C &= \begin{bmatrix} a_{11}\text{vec}B \\ a_{1n}\text{vec}B \\ \vdots \\ a_{m1}\text{vec}B \\ a_{mn}\text{vec}B \end{bmatrix} \\ &= \text{vec}(a_{11}\text{vec}B, \dots, a_{1n}\text{vec}B, \dots, a_{m1}\text{vec}B, \dots, a_{mn}\text{vec}B) \\ &= \text{vec}\{(\text{vec}B)(\text{vec}A')'\}, \end{aligned}$$

but also

$$\text{vecb}_r C = \text{vec}(a_{11}B, \dots, a_{1n}B, \dots, a_{m1}B, \dots, a_{mn}B) = \text{vec}\{(\text{vec}A')' \otimes B\}.$$

As a second result we have that, if  $C = (\text{vec}B)(\text{vec}A)'$ , where  $B$  is elementwise partitioned, then

$$X = \text{vec}B \otimes \text{vec}A \tag{24}$$

is a solution for the equation  $\text{vec}X = \text{vecb}_r C$ . This can be seen as follows:

$$C = (\text{vec}B)(\text{vec}A)' = \begin{bmatrix} b_{11}(\text{vec}A)' \\ b_{m1}(\text{vec}A)' \\ \vdots \\ b_{1n}(\text{vec}A)' \\ b_{mn}(\text{vec}A)' \end{bmatrix},$$

say. Then

$$\text{vecb}_r C = \begin{bmatrix} b_{11}\text{vec}A \\ b_{m1}\text{vec}A \\ \vdots \\ b_{1n}\text{vec}A \\ b_{mn}\text{vec}A \end{bmatrix} = \text{vec}B \otimes \text{vec}A = \text{vec}(\text{vec}B \otimes \text{vec}A).$$

#### 4 APPLICATIONS

In this section we will give some examples where the tilde operator and both block Kronecker products play an important role. Various results concerning the block Kronecker product will be used.



#### 4.1 Approximation of a large matrix

In the context of covariance structure analysis, it can be computationally attractive to approximate a  $(k^2 \times k^2)$  matrix  $W$  by  $A \otimes A$  (see Koning, Neudecker and Wansbeek [3]). The criterion to be minimized is

$$\text{tr}[A \otimes A - W]^2, \quad (25)$$

where  $A$  is a symmetric  $(k \times k)$  matrix as yet to be determined and  $W$  is a symmetric  $(k^2 \times k^2)$  matrix of fourth-order moments, hence  $K_{kk}W = WK_{kk} = W$ . Denoting the criterion function by  $k(a)$ , with  $a \equiv \text{vec}A$ , we have

$$\begin{aligned} k(a) &= \text{tr}[(A \otimes A) - W]^2 = (a \otimes a)' B' B (a \otimes a) - 2(a \otimes a)' B' W + \text{tr}W^2 \\ &= (a'a)^2 - 2a' \tilde{W} a + \text{tr}W^2 = (\text{tr}A^2)^2 - 2a' \tilde{W} a + \text{tr}W^2 \end{aligned} \quad (26)$$

with  $B = I_k \otimes K_{kk} \otimes I_k$ . For this function we derive the first differential. In the derivation, the matrix  $D$  is the duplication matrix (see Magnus and Neudecker [5]), and  $\psi(A)$  is the vector with nonduplicated elements of  $\text{vec}A$ . The first differential of  $k(a)$  is

$$dk(A) = 4(\text{tr}A^2)a'da - 4a'\tilde{W}da = 4(\text{tr}A^2)a'Dd\psi(A) - 4a'\tilde{W}Dd\psi(A),$$

since

$$D\psi(A) = \text{vec}A = a.$$

The first-derivative is now

$$\frac{\partial k(A)}{\partial \psi(A)} = 4(\text{tr}A^2)D'\text{vec}A - 4D'\tilde{W}\text{vec}A,$$

from which we obtain the first order condition

$$(\text{tr}A^2)D'\text{vec}A = D'\tilde{W}\text{vec}A,$$

which leads to

$$(\text{tr}A^2)\text{vec}A = \tilde{W}\text{vec}A. \quad (27)$$

The first-order condition of the minimization problem (25) thus takes the



form of an eigenvalue problem in terms of  $\tilde{W}$ . It is easily seen that the vector  $a^*$  minimizing (26) must be proportional to the eigenvector corresponding to the largest eigenvalue. In fact, if  $\lambda^*$  is the largest eigenvalue of  $\tilde{W}$  and  $x^*$  is the corresponding eigenvector,  $a^* = \sqrt{\lambda^*} x^*$  minimizes (26).

## 4.2 $k$ -factorial covariance structures

In the estimation of  $k$ -factorial covariance structures, the distance between the  $l \times l$  sample covariance matrix  $S$  and the population covariance matrix  $\Omega$  is minimized. The matrix  $\Omega$  is restricted to have the form

$$\Omega = \Sigma_1 \otimes \Sigma_2 \otimes \dots \otimes \Sigma_k,$$

with  $\Sigma_i$  of order  $(l_i \times l_i)$ . Of course,  $\prod_{i=1}^k l_i = l$ . A general form of the criterion function used in this problem is

$$q(\theta) = \text{tr}[(S - \Omega)W^{-1}]^2, \quad (28)$$

with  $q$  the function to be minimized,  $\theta$  the vector of parameters to be estimated (i.e., the elements of  $\Sigma_i$ ,  $i = 1, \dots, k$ ) and  $W$  is a weighting matrix.

To derive estimators for  $\Sigma_i$ , some additional notation is needed. Define

$$W_i \equiv C_i W C_i',$$

with  $C_i \equiv I_{l_1 \times \dots \times l_{i-1}} \otimes K_{l_{i+1} \times \dots \times l_k, l_i}$ . This matrix  $C_i$  is a generalization of the commutation matrix and is discussed extensively in Kapteyn, Neudecker and Wansbeek [2]. The matrices  $S_i$  and  $\Omega_i$  are defined analogously. Note that

$$\Omega_i \equiv C_i \Omega C_i' = (\Sigma_1 \otimes \dots \otimes \Sigma_{i-1} \otimes \Sigma_{i+1} \otimes \dots \otimes \Sigma_k) \otimes \Sigma_i \equiv \Sigma^i \otimes \Sigma_i,$$

with  $\Sigma^i$  implicitly defined. Furthermore, define  $l^i \equiv \prod_{j \neq i}^k l_j$ ,  $\sigma_i \equiv \text{vec} \Sigma_i$  and  $\sigma^i \equiv \text{vec} \Sigma^i$ . Finally, we will use the matrix  $B_i$ , which is given by  $B_i = I_{l^i} \otimes K_{l^i, l_i} \otimes I_{l_i}$ . Now we are in a position to evaluate the criterion function:

$$\begin{aligned} q(\theta) &= \text{tr}[(S - \Omega)W^{-1}]^2 = \text{tr}[(S_i - \Omega_i)W_i^{-1}]^2 \\ &= [\text{vec}(S_i - \Omega_i)]' (W_i \otimes W_i)^{-1} [\text{vec}(S_i - \Omega_i)] \\ &= [\text{vec}(S_i - \Omega_i)]' B_i' B_i (W_i \otimes W_i)^{-1} B_i' B_i [\text{vec}(S_i - \Omega_i)] \\ &= [\text{vec} S_i - X_i \sigma_i]' (W_i \otimes W_i)^{-1} [\text{vec} S_i - X_i \sigma_i], \end{aligned} \quad (28')$$



with  $X_i \equiv \sigma_i^i \otimes I_{l_i} \otimes I_{l_i}$ . In this derivation, we have used equation (11) and the property of the tilde operator in equation (7). An estimator for  $\sigma_i$  is thus given by

$$\hat{\sigma}_i = [X_i' (W_i \pi W_i)^{-1} X_i]^{-1} X_i' (W_i \pi W_i)^{-1} \text{vec} \tilde{S}_i. \quad (29)$$

A more elaborate discussion of this model can be found in Verhees [10] and Wansbeek and Verhees [12].

### 4.3 The Moore-Penrose inverse of a design matrix in balanced ANOVA

Consider the usual linear balanced ANOVA model with fixed effects:

$$y_{ij} = \alpha + \beta_i + \gamma_j + \varepsilon_{ij}, \quad \begin{array}{l} i = 1, \dots, n \\ j = 1, \dots, m \end{array} \quad (30)$$

with  $\varepsilon_{ij}$  a random disturbance term. In matrix format, this can be written as

$$\begin{aligned} y &= (\iota_m \otimes \iota_n \ I_m \otimes \iota_n \ \iota_m \otimes I_n) \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} + \varepsilon \\ &\equiv X\theta + \varepsilon. \end{aligned} \quad (31)$$

Estimation of  $\theta$  requires the Moore-Penrose inverse of the design matrix  $X$ . It is convenient, however, to consider a larger matrix  $\bar{X}$ :

$$\bar{X} \equiv (I_m \otimes I_n \ \iota_m \otimes I_n \ I_m \otimes \iota_n \ \iota_m \otimes \iota_n).$$

If we define

$$\bar{X}_m \equiv (I_m, \ \iota_m),$$

then

$$\bar{X} = (I_m, \ \iota_m) \square (I_n, \ \iota_n) = \bar{X}_m \square \bar{X}_n. \quad (32)$$

The Moore-Penrose inverse is derived by utilizing a singular value decomposition of  $\bar{X}$ . If  $\bar{X}$  has a singular value decomposition  $\bar{X} = UDV'$ , then the Moore-Penrose inverse is given by  $\bar{X}^+ = VD^{-1}U'$ . Of course, a singular value decomposition can serve other purposes besides deriving its Moore-Penrose inverse, like assessing the rank of  $X$  or to characterize estimable functions



of  $\theta$ . To derive a singular value decomposition of  $X$ , we need one preliminary result (see Wansbeek [11]). A singular value decomposition of  $X_m$  is given by

$$X_m = U_m D_m V'_m \equiv (G_m, m^{-1/2} I_m) \begin{bmatrix} I_{m-1} & 0 \\ 0 & (m+1)^{1/2} \end{bmatrix} \begin{bmatrix} G'_m & 0 \\ \phi_m I'_m & m \phi_m \end{bmatrix}, \quad (33)$$

with  $G_m$  an  $m \times (m-1)$  matrix such that  $G'_m G_m = I_{m-1}$  and  $G'_m I'_m = 0$ . The constant  $\phi_m$  is defined by  $\phi_m \equiv (m(m+1))^{-1/2}$ . Now we are in a position to derive a singular value decomposition of  $X$ :

$$\begin{aligned} X &= X_m \square X_n \\ &= (U_m D_m V'_m) \square (U_n D_n V'_n) \\ &= (U_m \square U_n) (D_m \square D_n) (V'_m \square V'_n)' \\ &\equiv U D V', \end{aligned} \quad (34)$$

which is evidently a singular value decomposition of  $X$ . In deriving this result we have used properties 6 and 7 from section 2.2. A singular value decomposition of  $X$  can now be obtained as follows. If we define  $V'_R$  to be the matrix  $V'$  with the first block of columns deleted (corresponding to  $X$  being  $X$  with the first block of columns deleted), it is easily established that

$$V'_R V_R = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & \alpha_1 I_{n-1} & 0 & 0 \\ 0 & 0 & \alpha_2 I_{m-1} & 0 \\ 0 & 0 & 0 & \alpha_3 \end{bmatrix}, \quad (35)$$

with  $\alpha_1 = m/(m+1)$ ,  $\alpha_2 = n/(n+1)$  and  $\alpha_3 = (m+n+mn)/\{(m+1)(n+1)\}$ . Now define  $Q$  by:

$$Q \equiv \begin{bmatrix} 0 & 0 & 0 \\ I_{n-1} & 0 & 0 \\ 0 & I_{m-1} & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

then it is easily verified that a singular value decomposition of  $X$  is

$$X = (UQ) (Q'D[V'_R V_R]^{1/2} Q) (Q'([V'_R V_R]^+)^{1/2} V'_R). \quad (36)$$

See Wansbeek [11] for more details. This approach can be generalized to more complicated models.



## REFERENCES

1. D.C. Hyland and E.G. Collins, Block Kronecker products and block norm matrices in large-scale systems analysis, *SIAM Journal of Matrix Analysis and Applications*, 10 (1989), pp. 18–29.
2. A. Kapteyn, H. Neudecker and T. Wansbeek, An approach to  $n$ -mode components analysis, *Psychometrika*, 51 (1986), pp. 269–275.
3. R.H. Koning, H. Neudecker and T. Wansbeek, A note on approximately efficient estimation of covariance structure models, manuscript, (1989).
4. J.R. Magnus and H. Neudecker, The commutation matrix: some properties and applications, *Annals of Statistics*, 7 (1979), pp. 381–394.
5. J.R. Magnus and H. Neudecker, The elimination matrix: some lemmas and applications, *SIAM Journal on Algebraic and Discrete Methods*, 1 (1980), pp. 422–449.
6. H. Neudecker and T. Wansbeek, Some results on commutation matrices, with statistical applications, *Canadian Journal of Statistics*, 11 (1983), pp. 221–231.
7. Singh, R.P., *Some generalizations in matrix differentiation with applications in multivariate analysis*, Ph.D. Dissertation, University of Windsor (1972)
8. D.S. Tracy and K.G. Jinadasa, Partitioned Kronecker products of matrices and applications, *Canadian Journal of Statistics*, 17 (1989), pp. 107–120.
9. D.S. Tracy and R.P. Singh, A new matrix product and its applications in partitioned matrix differentiation, *Statistica Neerlandica*, 26 (1972), pp. 143–157.
10. J. Verhees, *Econometric analysis of multidimensional models*, Ph.D. Dissertation, University of Groningen (1989).
11. T. Wansbeek, Singular value decomposition of design matrices in ANOVA with balanced data, manuscript (1989).
12. T. Wansbeek and J. Verhees, Models for multidimensional matrices in econometrics and psychometrics, in R. Coppi and S. Bolasco (eds.), *Multiway data analysis*, Amsterdam: North-Holland (1989).



Papers issued in these series from 1989 onwards: <sup>1)</sup>

297. Steven Brakman and Charles van Marrewijk, Transfer problems in a multi-country world. Endogenous, optimal and coordinated aid (1989).
298. C.L.J. van der Meer and S. Yamada, Productivity and income in intensive livestock raising in Japan and The Netherlands in 1983/84 (1989).
299. Maarten C.W. Janssen and Yao-Hua Tan, Why Friedman's non-monotonic reasoning defies Hempel's covering law model (1989).
- 300.\* Günter Knieps, Deregulation in Europe: telecommunications and transportation (1989).
301. J. Oosterhaven, A review of Dutch rent control and its farreaching consequences (1989).
302. Gerard J. van den Berg, The theory of nonstationary job search and an empirical analysis of unemployment benefits as a decreasing function of duration (1989).
303. Robert Lensink, A model of rural-urban conflict in developing countries (1989).
304. Jan Jacobs and Maarten Janssen, Coordinating unions, wages and employment (1989).
305. Herman Cesar, On testing the intertemporal substitution effects of government purchases with military spending data (1989).
306. Herman Cesar, Jakob de Haan and Jan Jacobs, Monetary targeting in the Netherlands. An application of cointegration tests (1989).
307. K.J. Alsem, P.S.H. Leeftang en J.C. Reuyl, Concurrentieverhoudingen op de advertentiemarkt. Deel 2: een verklarende analyse (1989).
308. Albert W. de Groot en Robert Lensink, Stabilisatieprogramma's in ontwikkelingslanden: de bruikbaarheid van multi-sectormodellen (1989).
309. Charles van Marrewijk and Casper G. de Vries, The customs union argument for a monetary union (1989).
310. Evert Jan Bakker, Possible effectiveness of rainfall insurance as a means to reduce rural household risk in India; a preliminary analysis (1989).
311. Charles van Marrewijk and Peter A.G. van Bergeijk, Uncertain trade (1989).
312. Ch.B. Blankart and G. Knieps, What can we learn from comparative institutional analysis? The case of telecommunications (1989).
313. Geert Ridder, Attrition in multi-wave panel data (1989).
314. Gerard J. van den Berg, The theory of nonstationary job search and an empirical analysis of unemployment benefits as a decreasing function of duration (1989).
315. D.J. Nicolai, Th.J.B.M. Postma, Strategische planning- en voorspellingsmethoden in het Nederlandse bedrijfsleven: een inventarisatie (1989).
316. Niels Hermes en Robert Lensink, Het plan-Brady: de beslissende doorbraak in het schuldenvraagstuk? (1989).
317. Harry Garretsen and Maarten C.W. Janssen, Two fallacies of composition in a Keynesian olig model (1989).
318. Ronald Boin en Jouke van Dijk, Volunteer labor supply in the Netherlands: a tobit approach (1989).
319. Harry Garretsen and Maarten C.W. Janssen, Indeterminacy in macroeconomics: on the method of Keynes' general theory (1989).

---

<sup>1)</sup> These research memoranda can be ordered at the Institute of Economic Research, University of Groningen, P.O. Box 800, 9700 AV Groningen.



320. Peter A.G. van Bergeijk, Trade and diplomacy: an extension of the gravity model in international trade theory (1989).
321. D.F. Broens en L.R.J. Westermann, Toepassingen van Splinefuncties (1989).
322. Cars Hommes, Periodic, quasi-periodic and chaotic dynamics in a simple piecewise linear non-walrasian macromodel (1989).
323. M.A. Allers and E.H. van Leeuwen, Relative Acceptance in world clothing trade: the penetration pattern of the newly industrializing economies (1989).
324. Robert Lensink and Peter A.G. van Bergeijk, Development aid requirements in the 1990s (1989).
325. J.C. van Ours and G. Ridder, An empirical analysis of vacancy durations and vacancy flows: cyclical variation and job requirements (1989).
- 326.\* S.K. Kuipers en H. van Ees, Over de ontwikkelingen binnen de macro-economie na de tweede wereldoorlog en haar consequenties voor de economische politiek (1989).
327. Hans van Ees, Unemployment equilibria, efficiency wages and kinked demand curves: the scope for economic policy (1989).
328. Gerard J. van den Berg, Search behaviour, transitions to nonparticipation and the duration of unemployment (1989).
329. Wietse Dol and Ton Steerneman, Asymptotically optimal bilinear and QR estimators for small area parameters (1989).
330. Maryse J. Brand, Measuring industrial marketing mix effectiveness; a review (1989).
331. Bas Bakker, On the economic effects of government debt (1989).
332. Lambert Schoonbeek, The size of the balanced growth rate in the dynamic Leontief model (1989).
333. Hans van Ees and Harry Garretsen, The right answers to the wrong question? An assessment of the microfoundations debate (1989).
334. Robert Lensink and Peter A.G. van Bergeijk, Access to the international capital market: the determinants of international bank lending (1989).
335. J.S. Cramer and G. Ridder, Pooling states in the multinomial logit model (1989).
336. H.W. Hoen and E.H. van Leeuwen, Upgrading and relative competitiveness in manufacturing trade: Eastern Europe versus the newly industrializing economies (1989).
337. Albert W. de Groot, Adjustment policies and the real exchange rate in Kenya from 1975 onwards (1989).
338. Willem G. Nijkamp, Ranking multi-attributed alternatives by dominance-related concepts: a survey (1989).
339. W. Voorhoeve, H.W.A. Dietzenbacher, Het model GRECON 89-A; september-voorspellingen 1990 (1989).
340. K.J. Alsem, P.S.H. Leeftang en J.C. Reuyl, Concurrentieverhoudingen op de arbeidsmarkt. Deel 3: Een voorspellende analyse (1989).
341. Evert Schoorl, The institutionalization of political economy: the case of the Netherlands (1989).
342. Steven Brakman and Charles van Marrewijk, Endogenous transfers, a note on Bhagwati, Brecher and Hatta (1989).
343. T. Stam and G. Sierksma, The 3-interchange graph and the symmetric traveling salesman polytope (1989).
344. Theo K. Dijkstra, A note on description versus prediction (1989).
345. Geert Ridder and Insan Tunali, Analysis of related durations: a semi-parametric approach. With an application to a study of child mortality in Malaysia (1989).
346. Gerard J. van den Berg, A structural dynamic analysis of job turnover and the costs associated with moving to another job (1989).
347. E. Schoorl en C. Schweigman, Voorwaardelijk gefinancierd economisch onderzoek aan de RUG, 1984-1988 (1989).



- 348. Pierre van der Eng, Growth and productivity change in Indonesian agriculture, 1880-1985 (1990).
- 349. I.P. van den Berg, Exponential stability and the river-phenomenon (1990).
- 350. Charles van Marrewijk, An allocation mechanism, a measure of risk aversion and an experiment (1990).
- 351. Ruud H. Koning, Heinz Neudecker, Tom Wansbeek, Block Kronecker products and the vecb operator (1990).